



Rabbit anti-Human 3'Sulfo Lewis X Antibody

Product #:	GKGB4RBIGG100
Clone name:	E2-B4
Product type:	Recombinant rabbit IgG monoclonal antibody
Lot number:	T2234A23
Source:	Cell culture expression in HEK293 cells
Purification:	Purified by Protein A or Protein G affinity chromatography
Immunogen:	3'Sulfo-Lewis X
Buffer:	Phosphate buffered saline pH7.5
Endotoxin:	<0.050 EU/mg as determined by LAL chromogenic endotoxin assay
Concentration:	1mg/ml
Vial size:	0.1mg

Target information:


This antibody recognises the 3'sulfo- β Gal linkage found on the tetrasaccharide carbohydrate Sialyl LewisX (sLeX), usually found attached to O-glycans on the surface of cells. 3'-Sulfo-Lewis X (3-SLeX) is a glycan epitope that has been reported to act as a ligand for the adhesion molecules E- and L-selectin. The synthesis of 3-SLeX is mediated by the enzyme galactose-3-O-sulfotransferase (GAL3ST2), which is expressed in cell lines with high metastatic potential.

Expression of 3-SLeX is therefore potentially relevant to the identification of tumours with poor prognosis and higher likelihood of progression and metastasis.

3-SLeX has been found to be expressed at high levels in the mucins from the lungs of Cystic Fibrosis patients. It has been shown that *Pseudomonas Aeruginosa* preferentially binds to this form of LeX, and that high levels of expression may be related to more serious infections and chronic colonisation of airway mucins in Cystic Fibrosis.

Studies of 3-SLeX expression and function have often been inhibited by the lack of any specific antibodies. GlykoGen have overcome these reagent challenges by using their proprietary immunisation methods to develop Clone B4, a recombinant monoclonal rabbit IgG antibody. The availability of this antibody will facilitate studies into 3-SLeX expression and function, both in normal tissues and in cancer studies.


The chemical structure of the 3-SLeX antigen is shown below



Applications:


Immunohistochemistry: This antibody may be used to stain frozen tissue sections. A dilution range of 40 ug/ml is suggested, but users should titrate the antibody for best results within their own systems.

The image below shows staining of fresh frozen human colon cancer tissue, demonstrating antibody binding to individual cells and to discrete stromal components.



Immunoblotting: This antibody may be used for immunoblotting of lipid extracts of cells. A dilution range of 10-50 ug/ml is suggested, but users should titrate the antibody for best results within their own systems. NB: Due to the glycan nature of the epitope recognised by this antibody, traditional protein Western Blotting cannot be used.

The image below shows lipid immunoblotting of a total lipid extract of LS180 cells using the B4 antibody at 20ug/ml (with Sigma A0545 Goat anti-rabbit IgG peroxidase).



ELISA: This antibody may be used in ELISA applications. A dilution range of 20-50 ug/ml is suggested, but users should titrate the antibody for best results within their own systems

Recommended secondary antibody: A suitable anti-rabbit IgG reagent such as product A0545 (Goat anti-rabbit IgG:HRP) or SAB3700963 (Donkey anti-rabbit IgG (H+L) F(ab')₂:FITC from Sigma is recommended

Shipping: The product is shipped at ambient temperature. Upon receipt store immediately as recommended below

Storage: Use a manual defrost freezer and avoid repeated freeze/thaw cycles. Store frozen in aliquots at -20°C for up to one year, or at 4°C for up to one week.

References

1. Yuen CT, Bezouska K, O'Brien J, et al. Sulfated blood group Lewisia. A superior oligosaccharide ligand for human E-selectin. *Journal of Biological Chemistry*. 1994;269(3):1595-1598. doi:10.1016/s0021-9258(17)42065-5
2. Green PJ, Tamatani T, Watanabe T, et al. High affinity binding of the leucocyte adhesion molecule L-selectin to 3'-sulphated-Lea and -Lex oligosaccharides and the predominance of sulphate in this interaction demonstrated by binding studies with a series of lipid-linked oligosaccharides. *Biochem Biophys Res Commun*. 1992;188(1). doi:10.1016/0006-291X(92)92376-9
3. Yuen CT, Stoll MS, Feizi T, et al. Novel Sulfated Ligands for the Cell Adhesion Molecule E-Selectin Revealed by the Neoglycolipid Technology among O-Linked Oligosaccharides on an Ovarian Cystadenoma Glycoprotein. *Biochemistry*. 1992;31(38):9126-9131. doi:10.1021/bi00153a003

For Research Use Only. Not to be used for therapeutic or diagnostic applications.

Glykogen Ltd, The James Hutton Institute, Craigiebuckler, Aberdeen

Available in Canada from...

MJS
BioLynx
INC.

1-888-593-5969 • www.biolynx.ca • tech@chromspec.com

